

## Oddball distractors demand attention: neural and behavioral responses to predictability in the flanker task

Abigail Noyce

Department of Psychological and Brain Sciences  
Boston University

Robert Sekuler

Volen Center for Complex Systems  
Brandeis University

### Abstract

Predictable and unpredictable distractors may have very different attentional effects. We adapted the Eriksen flanker task by manipulating the probability with which specific flankers occurred. Subjects reported the orientation of briefly-presented targets, while attempting to ignore four flanking items. Flankers had either standard (90% of trials) or oddball (10%) orientations. Congruent and incongruent configurations were equiprobable, as were target orientations. Oddball flanker orientations substantially enhanced the congruency effect: performance was best when the target was congruent with oddball flankers and worst when it was incongruent. From subjects' electroencephalogram, we extracted the visual mismatch negativity (vMMN) evoked by oddball flanker orientations. Individual differences in vMMN magnitude were predicted by two factors of temperament, orienting sensitivity and attentional control. Subjects appear to exploit distractor predictability to support active inhibition; oddballs disrupt this strategy. Despite subjects' attempts to ignore the flankers, processing of unexpected distractors strongly influences neural responses and behavioral performance.

The natural environment's predictable spatial and temporal regularities allow the human brain to operate in a predictive, feedforward mode (Bar, 2009). This ability to extend environmental structure to predict forthcoming stimuli facilitates many cognitive tasks, from identifying objects (Biederman, Mezzanotte, & Rabinowitz, 1982) to planning and executing behaviors (Maryott, Noyce, & Sekuler, 2011) to appropriately allocating attention (Posner, 1980). Most re-

search into predictive processing has considered the impact of regularities among a task's targets, a focus which is entirely understandable, as such regularities clearly facilitate cognitive performance. However, little work has been done on regularities among task-irrelevant distractors. Everyday experience suggests that, for example, it is easier to ignore a train whistle that blows at the same time every day than to ignore a one that occurs at random. Further, we know that attention plays two complementary roles in cognition. We direct attention to targets and we withdraw attention from, or perhaps actively inhibit, distractors (James, 1890). If predictable distractors facilitate such inhibition, then we should find enhanced attentional selectivity when distractors are predictable, and impaired selectivity when they are irregular. In order to investigate this proposition, we measured the behavioral and neural consequences of both predictable and oddball distractors.

Our study adapted the Eriksen flanker task (Eriksen & Eriksen, 1974), which entails interference between conflicting visual information. Specifically, the flanker task requires subjects to focus visual attention on a single target, such as a left-facing or right-facing chevron, while attempting to ignore surrounding items. The flanking distractors can either match or differ from the target, and the congruency between the flankers and the target influences the accuracy and reaction time with which subjects can report the target's orientation (e.g. Eriksen & Eriksen, 1974; White, Ratcliff, & Starns, 2011). Despite subjects' attempts to ignore the distractors, flankers that are incongruent with the central target interfere with processing, leading to reduced speed and accuracy on those trials (Schmidt & Dark, 1998). We modified the flanker task by manipulating the frequency with which different distractors appeared, creating predictable and oddball flankers.

To supplement our behavioral measures, we drew on event-related brain potentials (ERPs), which provide a direct measure of neural activity time-locked to specific events (Luck, 2005). Because we were interested in the effects that oddball flankers might have, we focused on a particular ERP component, the visual mismatch negativity (vMMN). This is an early, negative-going deflection in the ERP that occurs in response to occasional deviant elements within a sequence of visual stimuli that obey some regularity (Czigler, 2007; Pazo-Alvarez, Cadaveira, &

Amenedo, 2003). The vMMN is analogous to the well-established auditory mismatch negativity (MNN), which is theorized to be generated in auditory cortex when a predictive signal from prefrontal areas is disconfirmed by incoming sensory information (Garrido, Kilner, Stephan, & Friston, 2009; Wacongne, Changeux, & Dehaene, 2012). The auditory and visual MMNs do not depend on attention, and arise even when subjects are attending to stimuli in a different sensory modality (Näätänen, Paavilainen, Titinen, Jiang, & Alho, 1993; Stefanics, Kimura, & Czigler, 2011).


We measured the visual mismatch negativity in order to characterize the neural response to infrequent distractors, which subjects are actively attempting to ignore. We then assessed correlations between the neural responses, the behavioral effects of the frequent and infrequent distractors, and individual differences in temperament. We hypothesized that infrequent distractors would evoke a visual mismatch negativity, and that they would lead to an outside flanker congruency effect, due to the difficulty of suppressing attention to unpredictable flankers.

An area of growing interest among cognitive neuroscientists is the nature and origins of individual differences in neural activity and behavioral effects. One potential source of such differences is variability in emotional, motor, and attentional reactivity (Kagan, 2003; Rothbart, 2007). Such variability may predict the degree to which occasional oddball flankers disrupt subjects' attentional selectivity. We thus hypothesized that differences in temperament would predict differences in the extent to which oddball flankers enhance the flanker congruency effect. Further, the vMMN indicates that oddball stimuli are being processed, despite attention being directed away from them (Stefanics et al., 2011; Näätänen, Paavilainen, Rinne, & Alho, 2007), and we thus hypothesized that individual differences in temperament would predict the magnitude of the vMMN response to deviant distractors.

## Methods

### Subjects

Twenty members of the Brandeis University community (15 female, age range 18–21) participated in this study. All were right-handed (mean score on the revised Edinburgh Handedness



*Figure 1.* (A) The sequence of events within a trial. After fixation, the flanker stimulus was displayed for 50 ms, after which participants had two seconds to report the orientation of the center chevron. (B) Diagram of the two-by-two trial design. One factor, **flanker orientation**, governed the orientation of the four flanker chevrons; the second, **congruency**, governed the relationship between the central target and its flankers. Whether the **Standard** flanker orientation was left or right was counterbalanced within subjects. 90% of trials incorporated the **Standard** flankers (45% **Congruent** and 45% **Incongruent**); 10% of trials incorporated the **Oddball** flankers (5% **Congruent** and 5% **Incongruent**).

Inventory 89.49, SD = 12.85). Two other subjects completed one experimental session but did not return for the second; their data were discarded. The sample size of 20 was selected before data collection began, after considering (1) the magnitude of the interaction between flanker orientation and congruency, as seen in pilot data, (2) the amplitude of the visual mismatch negativity, as estimated from previous literature, and (3) the correlation strengths we wanted to be able to detect.

### Experimental task

We developed a modified Eriksen flankers task using chevron stimuli (Eriksen & Eriksen, 1974). The basic trial structure is shown schematically in Figure 1A. On each trial, subjects were presented with an array of five chevrons that were displayed for 50 ms and were not masked

upon offset. They were instructed to report whether the central chevron was pointing to the left or to the right. We will refer to this central chevron as the *target*, and the two chevrons on each side of it as the *flankers*. The four flankers were always consistently oriented, and the central chevron's orientation was equiprobably congruent or incongruent with its flankers. After a subject's response, a fixation cross was displayed for an inter-trial interval of 1000 ms before the next trial display appeared.

Subjects viewed the display from a distance of approximately 57 cm, and were instructed to maintain fixation on a central cross. Each chevron subtended approximately 1.40° visual angle, and the full array extended to an eccentricity of 4.67° to the left and right of the fixation point..

In order to maintain more-consistent error levels across subjects and conditions, subjects received feedback about their performance after every thirty trials (after Hajcak & Foti, 2008). If the subject had responded correctly on between 75% and 90% of those trials, the feedback was "You're doing great!" If accuracy was lower than 75%, the feedback instructed the subject to increase their accuracy; if it was above 90%, the feedback instructed the subject to respond more quickly.

Trials were randomly distributed among four conditions in a two-by-two design, as shown in Figure 1B. The first factor governed the orientation of the four flanker chevrons. On ninety percent of trials, the flanker chevrons had one orientation (the **Standard** orientation) and on ten percent of trials they had the other (the **Oddball** orientation). The second factor governed the relationship between the central target and the flankers. On half of trials, the target was **Congruent** with the flankers, and on half it was **Incongruent**. Left-facing and right-facing targets were equally frequent, and the orientations comprising **Standard** and **Oddball** flankers were counter-balanced within subjects.

On approximately 13% of trials, a burst of auditory white noise was presented after either stimulus presentation or after response, to elicit startle reactions. Startle blink was measured via electromyography; however, we observed no effects of timing or trial type on startle blink magnitude and those data are not presented here.

### **Procedures and Analyses**

Each recording block comprised 510 trials, with the first thirty discarded as practice. Each subject completed two blocks with left-facing flankers as the **Standard** orientation and two with right-facing flankers. These four blocks were completed in two separate recording sessions; the order of blocks was counterbalanced across subjects. By the end of the experiment, each subject had completed 2,040 trials, 1,920 of which were included for analysis.

Subjects filled out an anonymous questionnaire after each recording session, confirming that they got reasonable amounts of sleep, were not under the influence of any psychoactive substances, and had no medical history, such as a head injury or neurological diagnosis, that would lead us to exclude their data.

**Behavioral measures.** Subjects' reaction times and responses were recorded from each trial and analyzed. We computed accuracy and median reaction time for each of **Congruent Standard**, **Incongruent Standard**, **Congruent Oddball**, and **Incongruent Oddball** conditions as well as Vincentile reaction times and accuracy for each condition (Vincent, 1912; Ratcliff, 1979).

After the end of their final experimental session, subjects completed the Adult Temperament Questionnaire Short Form (ATQ). This instrument has 77 items which form several self-report scales describing temperament factors (Evans & Rothbart, 2007). We selected two factors a priori for testing: Attentional Control and Orienting Sensitivity. Attentional Control refers to the capacity to focus attention, and to shift attention as desired. "It's often hard for me to alternate between two different tasks," is an example of a reverse-scored Attentional Control item. Orienting Sensitivity refers to awareness of low-intensity environmental and self-generated stimuli and experiences. "I often notice visual details in the environment," is an example of an Orienting Sensitivity item. We hypothesized that the Attentional Control would account for some variability in people's task performance, and that both would relate to variability in the ERPs elicited by **Oddball** and **Standard** stimuli.

**EEG recording and analyses.** A high-density EEG system (Electrical Geodesics, Inc., Eugene, OR) with 129 electrodes sampled scalp electroencephalographic signals at 250 Hz using a high-impedance amplifier. Signals were recorded for later, off-line analysis. At the start of each

experimental session, all channels were adjusted for scalp impedance below 50 k $\Omega$  impedance; after one experimental block, channels impedances were measured and, if needed, returned to at most 50 k $\Omega$  scalp impedance before the subject completed the session.

After recording, EEG data were preprocessed using the EEGLAB Matlab toolbox (Delorme & Makeig, 2004). Continuous EEG signals were bandpass filtered to between 0.25 and 100 Hz using a first-order Butterworth filter. A 60 Hz notch filter was also applied to the continuous data, to reduce line electrical noise. The data were then broken into epochs that were time-locked to stimulus onset and lasting from 200 ms before stimulus onset to 500 ms after. Epochs containing muscle artifacts, eye movements, and bad channels were identified by visual inspection and rejected. Independent components analysis allowed us to isolate eye blink activity, which was subtracted from the data. Data were again visually inspected for artifacts not corrected by the previous two processes. Finally, data were re-referenced to the average voltage, and averaged across trials and sessions to create a subject average ERP for each condition.

To compare ERPs evoked by trials with Oddball flankers to those evoked by trials with Standard flankers, we used a data-driven, non-parametric clustering approach to select time windows and electrodes for analysis (Maris & Oostenveld, 2007). The FieldTrip toolbox includes software implementing this approach (Oostenveld, Fries, Maris, & Schoffelen, 2011). It first quantifies the difference between two conditions at each electrode, at each time point, using Student's  $t$ . Then, clusters of time- and/or space-adjacent electrodes with  $|t| > t_{\text{criterion}}$  are identified. Criterion  $t$ -values were selected by the experimenters after considering several factors, including the degrees of freedom of the comparison, the magnitude of the difference between the conditions, and the degree of spatial and temporal specificity desired.

To test significance, the  $t$ -scores of each cluster's member electrodes and time-points were summed, giving a cluster score that reflected both the extent of the cluster (in space and time) and the magnitude of the difference between the conditions at those electrodes and time points. A reference distribution of test statistics was generated by randomly permuting the data across the two conditions being compared, computing such scores for each resulting cluster, and taking the largest such cluster score on each of 1,000 permutations. Where cluster-wise  $p$ -values are

reported, they have been derived by comparing the empirically-obtained cluster score to such a reference distribution. Grand average ERPs were created by averaging across subjects and across the electrodes identified as part of the cluster.

## Results

### Behavioral measures




Figure 2. Reaction time and proportion correct for each decile of Standard (blue) and Oddball (green) trials, when the flankers are Congruent (solid lines) and Incongruent (dashed lines).

Figure 2 shows the effects of congruency and flanker orientation on subjects' speed and accuracy in the flanker task. On trials in which the flankers had the **Standard** orientation (presented in blue in Figure 2), subjects were faster and more accurate on **Congruent** trials (proportion correct 0.93 [0.91, 0.95]; median correct reaction time 356 ms [342, 370]) than on **Incongruent** trials (proportion correct 0.86 [0.84, 0.88]; median correct reaction time 378 ms [362, 394]), with the largest accuracy differences occurring on trials where subjects responded quickly. This effect was exaggerated when the flankers had the **Oddball** orientation (presented in green in Figure 2). On **Oddball Congruent** trials, subjects were faster and more accurate (proportion correct 0.95 [0.93, 0.97]; median correct reaction time 343 ms [329, 357]) than on **Standard Congruent** trials, and on **Oddball Incongruent** trials, subjects were slowest and least accurate (proportion correct 0.59 [0.52, 0.66]; median correct reaction time 427 ms [406, 448]).

The interaction effect (difference between congruency effect on **Oddball** trials and the




congruency effect on **Standard** trials) was 0.23 proportion correct [0.24, 0.36] and -20 ms [-28, -13]. These results were confirmed by two-way repeated measures ANOVAs with factors **congruency** and **flanker orientation**. There was a main effect of **congruency** on proportion correct ( $F(1,19) = 94.211, p < .0001$ ), a main effect of **flanker orientation** on proportion correct ( $F(1,19) = 80.019, p < .0001$ ), and a **congruency**  $\times$  **flanker orientation** interaction ( $F(1,19) = 101.809, p < .0001$ ). There was a significant main effect of **congruency** on median correct reaction time ( $F(1,19) = 92.244, p < .0001$ ), no main effect of **flanker orientation** on median correct reaction time ( $F(1,19) = 0.026, p = .8747$ ), but a significant **congruency**  $\times$  **flanker orientation** interaction ( $F(1,19) = 30.917, p < .0001$ ).

On average, subjects scored near the center of the temperament self-report range from the ATQ. The mean Perceptual Sensitivity score was 5.00 (SD = 0.86), and the mean Attentional Control score was 4.03 (SD = 0.91). There was an unconvincing negative correlation between Attentional Control and proportion correct ( $r = -0.291 [-.650, .174]$ ).

## ERPs

Figure 3 illustrates the neural activity that accompanied the presentation of **Standard** and **Oddball** flanker directions. A data-driven clustering and permutation analysis identified the cluster of electrodes that best captured ( $p < .001$ ) the difference in scalp voltage topographies between **Standard** and **Oddball** flankers. This cluster was derived with  $t_{\text{criterion}} = 3.883$ , the critical  $t$ -value at  $\alpha = .001$  and  $df = 19$ . The cluster was composed of 16 posterior electrodes that were more negative-going on **Oddball** than on **Standard** trials during an interval from 180–320 ms after stimulus onset. Figure 3A depicts the locations of the electrodes making up the cluster and the distribution of  $t$ -scores between ERPs to the two flanker orientations during that time window. The traces in Figure 3B show ERPs at the cluster, time locked to stimulus onset, to **Standard** (blue) and **Oddball** (green) flankers. Figure 3C shows the difference between the two traces in Figure 3B. Negative values reflect **Oddball**-evoked ERPs that were more negative-going than **Standard**-evoked ERPs.

We tested whether vMMN magnitude reflected subjects' susceptibility to interference




*Figure 3.* ERPs to **Standard** and **Oddball** flanker orientations at a cluster of posterior electrodes. A: Topographical plot showing the locations of the 16 electrodes making up the cluster, and the distribution across the scalp of  $t$ -values at 180–320 ms after stimulus onset. B: Traces show ERPs time locked to stimulus onset for **Standard** (shown in blue) and **Oddball** (green) flankers, at the cluster of electrodes shown in panel A. Error bars are within-subject s.e.m. C: Trace depicts the difference between the two traces shown in panel B. Negative values occurred when ERPs to **Oddball** flankers were more negative-going than those to **Standard** flankers.

from unexpected flanker orientations. To account for differences in neural function and anatomy that are unrelated to such susceptibility, we regressed out each subject's N2 magnitude to **Standard** flankers. N2 magnitude was operationalized as each subject's mean amplitude of the ERP from 180–220 ms after stimulus onset on **Standard** trials.

The residual MMN values were correlated with proportion correct on **Oddball Congruent** trials ( $r = -0.448$  [.006, .743]), such that those subjects who were most accurate on these trials had larger (negative) MMNs. However, residual MMN magnitude did not correlate significantly with proportion correct on **Oddball Incongruent** trials, ( $r = 0.194$  [-.273, .586]), nor did it correlate with any reaction time measures ( $|r| < .250$  for all relationships).

Figure 4 illustrates the relationship between temperament and vMMN magnitude. Resid-



*Figure 4.* Two self-report temperament factors, attentional control and orienting sensitivity, are significantly correlated with individual differences in vMMN magnitude. Higher scores on attentional control predict larger (more negative) vMMNs; higher scores on orienting sensitivity predict smaller vMMNs.

ual vMMN values were marginally negatively correlated with Attentional Control score ( $r = -0.439, [-.738, .005]$ ), suggesting that subjects who scored higher on Attentional Control tended to have larger (negative) MMNs, and positively correlated with Orienting Sensitivity score ( $r = 0.464 [.026, .978]$ ), such that subjects who scored higher on Orienting Sensitivity tended to have smaller vMMNs. Figure 4 illustrates the relationship between these three measures. Combining Attentional Control and Orienting Sensitivity in a multiple linear regression significantly predicted N2-corrected vMMN magnitude ( $R^2 = 0.35, p = .026$ ).

### Discussion

We modified the Eriksen flanker task by manipulating flanker frequency to create Standard and Oddball flanker directions. We replicated the usual flanker congruency effect: subjects were faster and more accurate when flankers were congruent with the target. As we hypothesized, this effect was strongly modulated by the predictability of the flankers. On trials with Oddball flankers that were congruent with the target, subjects were fastest and most accurate, while on trials with Oddball flankers that were incongruent with the target, subjects were worst.

This result is consistent with previous work demonstrating that unexpected or poorly-predicted events capture attention in associative learning tasks (Wills, Lavric, Croft, & Hodg-

son, 2007), and that deviant events in an unattended location or sensory modality impair target detection (Ljungberg & Parmentier, 2012; Nössl, Marsh, & Sörqvist, 2012). While there have been some prior investigations into the effects of predictability in the flankers task, these have been restricted to manipulations of the frequency of incongruent trials (e.g. Gratton, Coles, & Donchin, 1992; Purmann, Badde, Luna-Rodriguez, & Wendt, 2011), and the frequency of particular flanker/target combinations (Lehle & Hübner, 2008; Wendt & Luna-Rodriguez, 2009). Those investigations have suggested that when incongruent trials are likely, people adjust their level of cognitive control and are less influenced by flankers (Gratton et al., 1992).

Several mechanisms for that adjustment have been proposed. One is that the presence of conflict on an incongruent trial leads to an activation of cognitive control mechanisms, such that reaction times on subsequent trials are increased and the flanker congruency effect is decreased (Gratton et al., 1992). This effect occurs on both a trial-by-trial timescale and a more global timescale, such that blocks with frequent incongruent trials have reduced flanker congruency effects relative to blocks with infrequent incongruent trials. Another proposed mechanism is that short-term priming effects facilitate responses to repeated stimuli (Mayr, Awh, & Laurey, 2003; Wendt & Luna-Rodriguez, 2009) and repeated responses (Mordkoff, 1996). Neither of these can entirely explain our behavioral results. We found a reduced flanker congruency effect of Standard (frequently-occurring) flankers and an enhanced effect of Oddball flankers, despite holding the probability of incongruent trials stable at fifty percent. Our data thus support active inhibition of predictable flankers. Previous work in a related Stroop task has shown that explicit cues as to the nature of the upcoming distractor can dramatically reduce interference, due to recruitment of an active inhibition process (Chao, 2011). Here, subjects may develop a template for probable flanker identities based on previous trials. On a trial whose flankers match that template, subjects are more successful at inhibiting the flankers; on a trial whose flankers differ from the template, subjects are markedly less successful.

Our results also confirmed our hypothesis that Oddball flanker orientations would evoke a visual mismatch negativity. This ERP component is thought to reflect a potentiation of the neural response to stimuli that do not match a predictive feedback signal sent from higher cortical areas

to the sensory cortices (Garrido et al., 2009; Wacongne et al., 2012). The presence of a vMMN to the Oddball flankers confirm that subjects' brains are sensitive to the regularity governing flanker direction. While previous work establishing the mismatch negativity in the visual domain has not demonstrated interactions between deviants in the unattended region and behavioral performance, here we've shown that human sensitivity to unexpected events elicits a reliable neural signature and interferes with spatially-precise attentional selectivity.

We also found, as hypothesized, that subjects' self-reported temperament could predict their neural sensitivity to Oddball flankers. Two factors—Attentional Control and Orienting Sensitivity—together predicted about a third of the variance in vMMN magnitude. High Attentional Control scores may reflect proficiency at predictive inhibition of distractors, leading to larger differences between the neural responses to Standards and to Oddballs. Similarly, high Orienting Sensitivity may lead to strong sensory responses to both Oddball and Standard flanker stimuli, reducing such differences. Attentional Control scores have previously been linked to increased conflict-related negativity at anterior electrodes (Kanske & Kotz, 2012), and to alpha-band activity in the parietal lobe (Alfonso, Miquel, Xavier, & Blanca, 2013), further supporting the link between self-reported temperament and neural markers of attention.

Neither temperament nor vMMN magnitude was related to individual differences in performance on the flanker task. It may be the case that the magnitude of the mismatch signal generated in visual cortex, as detectable by scalp EEG, is unrelated to the attentional capture elicited by such a mismatch. The null relationship between self-reported temperament and behavior is more surprising, especially given the relationship between temperament and vMMN. It is possible that the feedback subjects were given, which encouraged them to maintain their performance within a particular range of accuracy, has masked any underlying differences in behavior. Further work investigating this system is called for.

Using a flanker task with predictable and oddball flanking distractors, we found that oddball flankers enhance the flanker congruency effect and elicit a visual mismatch negativity. Further, individual differences in vMMN magnitude correlate with individual differences in temperament, such that temperament accounts for roughly a third of vMMN variation. Although a sub-

stantial body of previous work has demonstrated the fallibility of human attentional selection, these are the first results showing a dramatic interaction between the probability of a given distractor and its behavioral effects. The human brain leverages structure and predictability among non-target distractors in order to support attentional selection.

### References

- Alfonso, M.-R. R., Miquel, T.-F. E., Xavier, B., & Blanca, A.-S. S. (2013). Resting parietal electroencephalogram asymmetries and self-reported attentional control. *Clinical EEG and Neuroscience, 44*, 188–192. doi: 10.1177/1550059412465871
- Bar, M. (2009). The proactive brain: Memory for predictions. *Philosophical Transactions of the Royal Society B: Biological Sciences, 364*, 1235–1243. doi: 10.1098/rstb.2008.0310
- Biederman, I., Mezzanotte, R. J., & Rabinowitz, J. C. (1982). Scene perception: Detecting and judging objects undergoing relational violations. *Cognitive Psychology, 14*, 143–177.
- Chao, H.-F. F. (2011). Active inhibition of a distractor word: The distractor precue benefit in the Stroop color-naming task. *Journal of Experimental Psychology: Human Perception and Performance, 37*, 799–812. doi: 10.1037/a0022191
- Czigler, I. (2007). Visual mismatch negativity. *Journal of Psychophysiology, 21*, 224–230. doi: 10.1027/0269-8803.21.34.224
- Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods, 134*, 9–21.
- Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & Psychophysics, 16*, 143–149.
- Evans, D. E., & Rothbart, M. K. (2007). Developing a model for adult temperament. *Journal of Research in Personality, 41*, 868–888. doi: 10.1016/j.jrp.2006.11.002
- Garrido, M. I., Kilner, J. M., Stephan, K. E., & Friston, K. J. (2009). The mismatch negativity: A review of underlying mechanisms. *Clinical Neurophysiology, 120*, 453–463. doi: 10.1016/j.clinph.2008.11.029
- Gratton, G., Coles, M. G., & Donchin, E. (1992). Optimizing the use of information: Strategic control of activation of responses. *Journal of Experimental Psychology: General, 121*, 480–506.
- Hajcak, G., & Foti, D. (2008). Errors are aversive: Defensive motivation and the error-related negativity. *Psychological Science, 19*, 103–108. doi: 10.1111/j.1467-9280.2008.02053.x
- James, W. (1890). *The principles of psychology*. New York, NY: Henry Holt.

- Kagan, J. (2003). Biology, context, and developmental inquiry. *Annual Review of Psychology*, *54*, 1-23. doi: 10.1146/annurev.psych.54.101601.145240
- Kanske, P., & Kotz, S. A. (2012). Effortful control, depression, and anxiety correlate with the influence of emotion on executive attentional control. *Biological Psychology*, *91*, 88–95. doi: 10.1016/j.biopsycho.2012.04.007
- Lehle, C., & Hübner, R. (2008). On-the-fly adaptation of selectivity in the flanker task. *Psychonomic Bulletin & Review*, *15*, 814–818. doi: 10.3758/PBR.15.4.814
- Ljungberg, J. K., & Parmentier, F. B. R. (2012). Cross-modal distraction by deviance: Functional similarities between the auditory and tactile modalities. *Experimental Psychology*, *59*, 355–363. doi: 10.1027/1618-3169/a000164
- Luck, S. J. (2005). *An introduction to the event-related potential technique*. Cambridge, MA: MIT Press.
- Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*, 177–190. doi: 10.1016/j.jneumeth.2007.03.024
- Maryott, J., Noyce, A., & Sekuler, R. (2011). Eye movements and imitation learning: Intentional disruption of expectation. *Journal of Vision*, *11*(1), 7:1–16.
- Mayr, U., Awh, E., & Laurey, P. (2003). Conflict adaptation effects in the absence of executive control. *Nature Neuroscience*, *6*, 450–452. doi: 10.1038/nn1051
- Mordkoff, J. T. (1996). Selective attention and internal constraints: There is more to the flanker effect than biased contingencies. In A. Kramer, G. D. Logan, & M. G. H. Coles (Eds.), *Converging operations in the study of visual selective attention* (p. 483-502). Washington, DC: American Psychological Association.
- Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic research of central auditory processing: A review. *Clinical Neurophysiology*, *118*, 2544–2590. doi: 10.1016/j.clinph.2007.04.026
- Näätänen, R., Paavilainen, P., Titinen, H., Jiang, D., & Alho, K. (1993). Attention and mismatch negativity. *Psychophysiology*, *30*, 436-450.
- Nöstl, A., Marsh, J. E., & Sörqvist, P. (2012). Expectations modulate the magnitude of attentional capture by auditory events. *PLoS One*, *7*, e48569. doi: 10.1371/journal.pone.0048569
- Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 156869. doi: 10.1155/2011/156869
- Pazo-Alvarez, P., Cadaveira, E., & Amenedo, E. (2003). MMN in the visual modality: A review. *Biological*

*Psychology*, 63, 199-236.

- Posner, M. I. (1980). Orienting of attention. *The Quarterly Journal of Experimental Psychology*, 32, 3-25. doi: 10.1080/00335558008248231
- Purmann, S., Badde, S., Luna-Rodriguez, A., & Wendt, M. (2011). Adaptation to frequent conflict in the Eriksen flanker task. *Journal of Psychophysiology*, 25, 50-59. doi: 10.1027/0269-8803/a000041
- Ratcliff, R. (1979). Group reaction time distributions and an analysis of distribution statistics. *Psychological Bulletin*, 86, 446-461.
- Rothbart, M. K. (2007). Temperament, development, and personality. *Current Directions in Psychological Science*, 16, 207-212. doi: 10.1111/j.1467-8721.2007.00505.x
- Schmidt, P. A., & Dark, V. J. (1998). Attentional processing of unattended flankers: Evidence for a failure of selective attention. *Perception & Psychophysics*, 60, 227-238.
- Stefanics, G., Kimura, M., & Czigler, I. (2011). Visual mismatch negativity reveals automatic detection of sequential regularity violation. *Frontiers in Human Neuroscience*, 5:46, 1-9. doi: 10.3389/fnhum.2011.00046
- Vincent, S. B. (1912). The function of the vibrissae in the behavior of the white rat. *Behavioral Monographs*, 1, 1-82.
- Wacongne, C., Changeux, J.-P. P., & Dehaene, S. (2012). A neuronal model of predictive coding accounting for the mismatch negativity. *Journal of Neuroscience*, 32, 3665-3678. doi: 10.1523/JNEUROSCI.5003-11.2012
- Wendt, M., & Luna-Rodriguez, A. (2009). Conflict-frequency affects flanker interference: Role of stimulus-ensemble-specific practice and flanker-response contingencies. *Experimental Psychology*, 56, 206-217. doi: 10.1027/1618-3169.56.3.206
- White, C. N., Ratcliff, R., & Starns, J. J. (2011). Diffusion models of the flanker task: Discrete versus gradual attentional selection. *Cognitive Psychology*, 63, 210-238. doi: 10.1016/j.cogpsych.2011.08.001
- Wills, A. J., Lavric, A., Croft, G. S., & Hodgson, T. L. (2007). Predictive learning, prediction errors, and attention: Evidence from event-related potentials and eye tracking. *Journal of Cognitive Neuroscience*, 19, 843-854. doi: 10.1162/jocn.2007.19.5.843